




1




6




11




2




7




12




3




8




13




4




9




14




5




10




15




I




A




C




E




M




H




O




N




B




D




J




G



F



K



L

$f'(x) > 0$  and  $f''(x) = 0$  everywhere.

$$f(x) = x^3 - x.$$

$f(x)$  has an inflection point at  $x = 1$  because  $f''(x)$  switches signs at  $x = 1$ .

÷

&

?

$f'(x) < 0$  and  $f''(x) = 0$  everywhere.

$f'''(x)$  switches signs at  $x = -1$ .

$f(x)$  has a vertical tangent line at  $x = -1$ .


★

@

!

$f'(x)$  has a jump discontinuity at  $x = 0$ .

$f(x)$  is always concave down because  $f'(x)$  is always decreasing.



=

□

+

$f(x)$  has a local minimum at  $x = -1$  because  $f'(x)$  switches signs from negative to positive there.  $f''(x)$  is constant.

$f'(0) = f''(0) = 0$  and  $f''(x)$  exists everywhere.


$f''(x)$  is undefined at  $x = 1$ .

△

#

♥

$f(x)$  and  $f'(x)$  are both periodic with period  $2\pi$ .



$$\int_0^2 f'(x) dx = 0.$$

∞

%

⇒